

= Research Note =

## Leukocyte count in follicular aspirate samples could be a prognostic biomarker of the developmental competence of human metaphase II oocytes

Takayuki ISHIKAWA<sup>1,2,†</sup>, Masakazu DOSHIDA<sup>1</sup>, Toshihiko KYOYA<sup>1</sup>,  
Taeka TAKISAWA<sup>1</sup>, Masahiro SAKURAI<sup>2</sup>, Yumi HOSHINO<sup>2</sup>,  
Kentaro TANEMURA<sup>2</sup>, Koichi KYONO<sup>1</sup> and Eimei SATO<sup>2</sup>

<sup>1</sup>Kyono ART Clinic, Sendai 980-0014, Japan

<sup>2</sup>Laboratory of Animal Reproduction, Graduate School of Agricultural Science, Tohoku University, Sendai 981-8555, Japan.

† Correspondence: tishikawa1227@nifty.com

### ABSTRACT

Multiple gestations markedly impact the health of children conceived via assisted reproductive technology (ART). The present study investigated the potential of leukocyte count in follicular aspirate (FA) samples as a follicular prognostic biomarker of the developmental competence of human metaphase II (MII) oocytes. FA samples were obtained from 27 ART patients during 27 natural menstrual cycles, Leukocytes were separated using density-gradient centrifugation and prepared the Diff-Quik stained smears. The 27 cycles were classified into 3 groups based on the developmental stage of the embryo (Group I, 7 oocytes (–); Group II, 6 unfertilized oocytes and 5 early cleavage embryos; Group III, 9 blastocysts) and evaluated estradiol, luteinizing hormone, and progesterone levels on the day of gonadotropin-releasing hormone agonist administration using enzyme immunoassay indicated that these levels did not correlate with developmental competence. The 27 cycles were divided into Categories A (19) and B (8) depending on the infiltrating leukocyte count in the follicular fluid (FF; less than  $2.5 \times 10^4$  cells/follicle, and the value or more, respectively). Further classification and analysis of the 19 cycles classified as Category A into 3 groups (Group I, 6 oocyte (–); Group II, 5 unfertilized oocytes and 3 early cleavage embryos; Group III, 5 blastocysts) showed no significant differences among the groups regarding infiltrating mononuclear leukocyte (IML) count but a significantly increased infiltrating polymorphonuclear leukocyte (IPL) count in Group III compared to Groups I and II. These findings suggest that IPL count in FF is associated with the blastocyst formation of human MII oocytes, and thus it can be a predictor of the developmental competence of MII oocytes.

**Key words:** follicular fluid, human metaphase II oocytes, *in vitro* fertilization, infiltrating polymorphonuclear leukocyte

### INTRODUCTION

Multiple gestations markedly impact the health and welfare of children conceived via assisted reproductive technology (ART). As twin pregnancies are associated with a higher risk of complications for both the mother and the fetus [1, 20, 21], reducing the incidence of

multiple gestations is often desirable. One reasonable approach to reduce the disadvantage is single-embryo transfer (SET). With the development of SET, non-invasive and objective methods of oocyte/embryo selection have been required.

As oocyte maturation is controlled in the follicular

Submitted: December 21, 2011

Accepted: December 30, 2013

Advance Publication in Website: December 31, 2013

microenvironment, age-related alteration of the microenvironment makes the human preovulatory oocyte susceptible to chromosomal disorders and cytoplasmic structural defects [8,9,25]. Because the follicular microenvironment is created by regulatory/metabolic factors present in the follicular fluid (FF) which is originated from follicular cells, detection of the existence of certain prognostic biomarkers in the FF might allow for prediction of oocyte viability. Higher levels of both follicular oxygen content [3,23,24] and vascular endothelial growth factor (VEGF) levels [14,16] in FF have been correlated with the developmental competence of human oocyte/embryo during *in vitro* fertilization (IVF) cycles. In addition, follicular concentrations of leptin [3,7], nitric oxide (NO) [3], high-density lipoprotein (HDL) cholesterol, HDL component proteins, and HDL micronutrients [5,6] have been reported to play important roles for in human oocyte/embryo development.

Although several studies have examined the presence of leukocytes in follicular aspirate (FA) samples obtained at oocyte retrieval in IVF cycles to determine their leukocyte populations [4,12,13,15,22], these studies used different methods of preparing the FA samples, estimating the leukocyte subset, and stimulating ovaries of patients, making the comparison of their results, which often differed, a difficult task. On the other hand, in their investigation of the leukocyte/erythrocyte ratio in FA samples and the leukocyte profile of both FA samples and peripheral blood in patients on the day of ovum pick up (OPU), Smith *et al.* (2005) concluded that leukocytes infiltrate the pre-ovulatory follicles [22]. Despite this finding, the origin of leukocytes remains unclear as a consequence of blood vessel damage during OPU.

Due to the mixed nature of the findings and the challenges faced in IVF research, no definitive predictor of oocyte viability has become widely used in clinical settings. This study evaluated infiltrating leukocyte count in FA samples as a prognostic biomarker of the developmental competence of human metaphase II (MII) oocytes.

The purpose of this experimental study is to evaluate the infiltrating leukocyte count in FF by examining FA samples obtained in natural cycles and to assess the potential of this variable as a predictor for developmental competence of human MII oocytes.

## MATERIALS AND METHODS

### Patients

FA samples were obtained during the 27 natural menstrual cycles of 27 patients with an unknown infertility factor undergoing an IVF cycle at the study site between December 2008 and October 2009 with the patients' informed consent.

### Semen preparation

Semen samples were prepared using Sil-Select Plus (FertiPro N.V., Beernem, Belgium) in accordance with the manufacturer's protocol. After gradient centrifugation, washing and swim-up preparation were performed. Routine semen analysis was then performed according to World Health Organization guidelines [26].

### Oocyte retrieval

Patients underwent transvaginal ultrasound and serum hormone analysis for estradiol (E<sub>2</sub>), luteinizing hormone (LH), and progesterone (P<sub>4</sub>) levels using enzyme immunoassay (Eclucys LH, E2 II and Progesterone II; Roche Diagnostics K.K, Tokyo, Japan). Hormone levels were measured from Day 13 to 16 of the menstrual cycle (Day 1: the first day of the menstrual cycle) on the day of gonadotropin-releasing hormone (GnRH) agonist (Buserecure; Fuji Pharma, Tokyo, Japan) administration. The schedule of oocyte retrieval was determined according to serum E<sub>2</sub>, P<sub>4</sub>, and LH levels. Final oocyte maturation was intranasally triggered with a GnRH agonist when the mean diameter of the dominant follicle was >18 mm under ultrasonography.

OPU was performed transvaginally at 36 h after GnRH agonist administration. For each cycle, the only dominant follicle was gently aspirated using a sterile syringe by a physician; no mechanical aspirators were used.

### Conventional *in vitro* fertilization and embryo culture

Conventional *in vitro* fertilization (cIVF) was performed with subsequent embryo culture [10]. In brief, all incubations were performed at 37°C in 6% CO<sub>2</sub>, 5% O<sub>2</sub>, and 89% N<sub>2</sub>. Oocyte-cumulus complexes (OCCs) were inseminated at a final concentration of  $1 \times 10^5$  motile sperm/ml at approximately 40 h post-trigger and incubated in Medi-Cult IVF medium (MediCult a/s, Jyllinge, Denmark) for 20 h. Inseminated oocytes were cultured in Global medium (Life Global, Guilford, CT, USA) from day 2 to 6 post-fertilization. Blastocysts were assessed by Gardner's criteria [11] and vitrified on day 5

**Table 1.** Preovulatory serum E<sub>2</sub>, LH and P<sub>4</sub> levels in patient on the day of GnRH-agonist administration

Group (n)	Oocyte(-)	Unfertilized/ early cleavage stage	Blastocyst	P value
	I (7)	I (11)	III (9)	
E <sub>2</sub> (pg/ml)	304.6 ± 89.8	306.0 ± 95.0	242.8 ± 53.8	NS
LH (mIU/ml)	7.8 ± 8.7	12.8 ± 6.7	8.2 ± 3.0	NS
P <sub>4</sub> (ng/ml)	0.42 ± 0.16	0.54 ± 0.25	0.63 ± 0.36	NS

E<sub>2</sub>: estradiol, LH; luteinizing hormone, P<sub>4</sub>; progesterone. Serum hormone levels are given as means ± SD. Non-repeated ANOVA method. NS: no statistical significance.

**Table 2.** Comparison of infiltrating leukocyte count in FF classified by developmental competence of MII oocytes

Developmental stage	Groups	Cycles (n)	IML count (10 <sup>2</sup> cells/follicle)	IPL count (10 <sup>2</sup> cells/follicle)
Oocyte(-)	I	6	28.5 ± 22.0 <sup>a</sup>	5.7 ± 7.2 <sup>c</sup>
Unfertilized	II	2	111.9 ± 22.1 <sup>b</sup>	8.4 ± 9.2 <sup>c</sup>
Early cleavage stage	II	6		
Blastocyst	III	5	159.9 ± 62.6 <sup>b</sup>	106.6 ± 30.7 <sup>d</sup>

IPL count; infiltrating polymorphonuclear leukocyte count, IML count; infiltrating mononuclear leukocyte count, oocyte (-); no oocyte was retrieved.

IML and IPL counts are expressed as means ± SD. Tukey-Kramer method; <sup>a-b, c-d</sup>P<0.01

or 6 using Cryotop (Kitazato Bio Pharma, Fuji, Japan).

### Leukocyte preparation

Leukocytes were separated from the FA samples using Polymorphrep (Axis-Shield, Oslo, Norway) in accordance with the manufacturer's protocol and centrifuged 2 times at 500 × g. After the leukocytes had been suspended in 100 μl of PBS, 1 μl suspensions were applied to prepare smears. The smears were fixed in methanol and stained with Diff-Quik (DQ) (CYSMEX, Kobe, Japan) in accordance with the manufacturer's protocol, with 2 separate smears prepared for each cycle. The stained smears were examined microscopically at × 400 or × 1,000 magnification and the leukocytes were classified as lymphocytes, neutrophils and eosinophils.

### Erythrocyte and leukocyte counts in FA samples

Based on the erythrocyte count in the FA samples and the blood cell populations in the peripheral blood of the patients on the day of OPU, the putative peripheral blood volume and peripheral leukocyte count in the FA samples were estimated. The infiltrating mononuclear leukocyte (IML) count in the FF was determined by subtracting the peripheral blood mononuclear leukocyte count from the mononuclear leukocyte count in the FA samples. The

infiltrating polymorphonuclear leukocyte (IPL) count in the FF was determined using a same method. As the FA samples contained varying amounts of flushing medium in each cycle, the leukocyte counts were expressed in terms of mean ± standard deviation (SD)/follicle.

### Grouping of cycles

The 27 cycles were classified into Category A (less than 2.5 × 10<sup>4</sup> cells/ follicle) and Category B (greater than 2.5 × 10<sup>4</sup> cells/follicle) depending on peripheral blood leukocyte count in the FA samples. The cycles that were classified as Category A were further classified into 3 groups based on the developmental stage of the embryo (Group I; no oocyte was retrieved, Group II; an unfertilized oocyte or an embryo at the early cleavage stage, and Group III; a blastocyst).

### Statistical analysis

Significant differences in hormone levels were evaluated by non-repeated measures ANOVA. On the other hand, infiltrating leukocyte counts in FF among the experimental groups were evaluated by Tukey-Kramer method. P values < 0.01 were considered to indicate statistical significance.

## RESULTS

### Patient age and oocyte retrieval

The mean patient age was  $33.6 \pm 3.1$  years (range: 27–38 years). OPU was performed for 27 cycles.

### cIVF and embryo culturing

In 20 of the 27 cycles, 1 oocyte was retrieved and cIVF was performed. As the results of cIVF and subsequent *in vitro* culturing, 14 fertilized oocytes (14/20, 70.0%) developed to the early cleavage stage and 9 blastocysts (9/14, 64.3%) were obtained on day 5 or 6. All blastocysts were classified as 6 Grade 4-blastocysts and 3 Grade 3-blastocysts, and vitrified them in liquid nitrogen.

### Serum hormone levels

The 27 cycles in which OPU was performed were classified into 3 groups based on the developmental stage of the embryos and statistically evaluated. As shown in Table 1, evaluation of  $E_2$ , LH, and  $P_4$  levels indicated that these levels did not correlate with the developmental competence.

### Peripheral blood leucocyte count in FA samples

Peripheral blood leucocyte count (peripheral blood volume) in the FA samples was less than  $2.5 \times 10^4$  cells/follicle ( $< 4 \mu\text{l}$ ) in 19 cycles (Category A) but greater than  $2.5 \times 10^4$  cells/follicle ( $> 6.8 \mu\text{l}$ ) in 8 cycles (Category B).

### Infiltrating leucocyte count in FA samples

The 19 cycles that were classified as Category A were further classified into 3 groups based on the developmental stage of the embryos. As shown in Table 2, although there were no significant differences between Group II and III with regard to IML count, there was a significant increase in the IPL count of Group III compared to that in Groups I and II. While IPL count ranged from 0.78 to  $1.59 \times 10^4$  cells/follicle in Group III cycles, it was less than  $0.29 \times 10^4$  cells/follicle in all Group I and II cycles. Due to the large number of peripheral blood leucocytes in the FA samples in all category B cases and the difficulty in determining IPL count on the stained smears, IPL count could not be determined accurately for this category.

## DISCUSSION

Oocyte maturation is known to be controlled a follicular microenvironment that is composed of follicular cells and regulatory/metabolic factors present

in the FF. Based on this knowledge, several investigators have investigated levels of  $O_2$ , VEGF, NO, leptin, HDL cholesterol, HDL component proteins, and HDL component micronutrients in FF as predictors of the developmental competence of human oocytes/embryos in IVF cycles [3,5–7,14,16,23,24]. In contrast, other investigators have focused on the presence of leucocytes in FF to determine leucocyte populations and the number of leucocytes in FA samples obtained at OPU in IVF cycles. However, the investigators have obtained mixed results due to their use of different methods and examination of different cycle characteristics patient populations, and could not determine the origin of leucocytes as a result of the presence of erythrocytes in the FA samples. Despite these challenges, in their investigation of the leucocyte/erythrocyte ratio in FA samples and the leucocyte profile of both FA samples and peripheral blood in patients on the day of OPU, Smith *et al.* (2005) concluded that leucocytes infiltrate the pre-ovulatory follicles [22].

In recognition of these findings and challenges, all FA samples examined in this study were obtained from patients in natural cycles, and contamination of the FA samples by peripheral blood due to blood vessel damage during OPU was considered. The putative peripheral blood volume was estimated based on erythrocyte count in both the FA samples and the peripheral blood, and the leucocyte count in the FA samples determined by subtracting the putative peripheral blood leucocyte count from the leucocyte count in the FA samples. Analysis of FA samples containing varying amounts of peripheral blood in which the peripheral blood leucocyte count was less than  $2.5 \times 10^4$  cells/follicle revealed a stronger correlation between blastocyst formation of MII oocytes and IPL count in FA samples obtained from follicles nurturing MII oocytes that developed to the blastocysts (Group III) compared with that of Groups I and II. On the other hand, the peripheral blood leucocyte count of category B exceeded  $2.5 \times 10^4$  cells/follicle, leading to difficulty in estimating the IPL count due to the large number of peripheral blood leucocytes on the DQ-stained smears.

As it involves the participation of both leucocytes and inflammatory mediators, ovulation, the terminal stage of oocyte maturation, shares several similarities with a local inflammatory reaction. In fact, previous studies have found that several inflammatory genes are expressed in preovulatory granulosa cells (GCs) and

cumulus cells (CCs) [17]. Specifically, the concentration of IL-8 (known as typical chemokine CXCL8) has been found to be higher in dominant follicles of the late follicular/ovulatory phase compared with those of the mid-follicular phase, and the capability of GCs to synthesize IL-8 to be up-regulated [18,19]. It is possible that the infiltrating polymorphonuclear leucocyte (IPML) populations of the FA samples examined in this study may have reflected the integrity of the microenvironment of the ovulatory follicle, specifically that of an environment that promoted oocyte maturation and ensured developmental competence of the human MII oocyte.

In conclusion, this is the first study to reveal that the IPL count of FA samples might be associated with the ability of human MII oocytes to develop to blastocysts. Although further studies are required using larger samples before IPL count can definitively defined a predictor in clinical settings and as a subset of IPL count populations in FA samples, the findings of this study indicate that IPL count may be a predictor of the developmental competence of human MII oocytes.

## REFERENCES

1. **Andersen AN, Goossens V, Ferraretti AP, Bhattacharya S, Felberbaum R, de Mouzon J.** The European IVF-monitoring (EM) Consortium for the European Society of Human Reproduction and Embryology (ESHRE). Assisted reproductive technology in Europe, 2004: results generated from European registers by ESHRE. *Hum. Reprod.* 2008; 23: 756–757.
2. **Antczak M, Van Blerkom.** Temporal and spatial aspects of fragmentation in early human embryos: possible effects on developmental competence and association with the differential elimination of regulatory proteins from polarized domains. *Hum. Reprod.* 1999; 14: 429–447.
3. **Barroso G, Barrionuevo M, Rao P, Graham L, Danforth, Huey S.** Vascular endothelial growth factor, nitric oxide, and leptin follicular fluid levels correlate negatively with embryo quality in IVF patients. *Fertil. Steril.* 1999; 72: 1024–1026.
4. **Beckmann MW, Polacek D, Seuag L, Schreiber JR.** Human ovarian granulosa cell culture: determination of blood cell contamination and evaluation of possible culture purification steps. *Fertil. Steril.* 1991 56: 881–887.
5. **Browne RW, Shelly WB, Bloom MS, Ocque AJ, Sandler JR, Huddleston HG, Fujimoto VY.** Distribution of high-density lipoprotein particle components in human follicular fluid and sera and their associations with embryo morphology parameters during IVF. *Hum. Reprod.* 2008; 23: 1884–1894.
6. **Browne RW, Bloom MS, Shelly WB, Ocque AJ, Sandler JR, Huddleston HG, Fujimoto VY.** Follicular fluid high-density lipoprotein-associated micronutrient levels are associated with embryo fragmentation during IVF. *J. Assist. Reprod. Genet.* 2009; 26: 557–560.
7. **Dorn C, Reinsberg J, Kupka M, van der Ven H, Schild RL.** Leptin, IGF-1 and IGFBP-3 concentrations in serum and follicular fluid of women undergoing *in vitro* fertilization. *Arch. Gynecol. Obstet.* 2003; 268: 187–193.
8. **Eichenlaub-Ritter U.** Genetics of oocyte ageing. *Maturitas.* 1998; 12: 143–169.
9. **Eichenlaub-Ritter U, Vogt E, Yin H, Gosden R.** Spindles, mitochondria and redox potential in ageing oocytes. *Reprod. Biomed. Online* 2004; 8: 45–58.
10. **Fauque P, Davy C, Jouannet P, Patrat C.** How to manage an accelerated embryo cohort in IVF. *Reprod. Biomed. Online* 2008; 16: 129–133.
11. **Gardner DK, Schoolcraft WB.** *In vitro* culture of human blastocysts. In Jansen R, Mortimer D (Ed.), *Toward reproductive certainty: fertility and genetics beyond 1999: the plenary proceedings of the 11th World Congress on In Vitro Fertilization and Human Reproductive Genetics*, The Parthenon Publishing Group, New York, 1999, pp. 378–388.
12. **Hill JA, Barbieri RL, Anderson DJ.** Detection of T8 (suppressor/cytotoxic) lymphocytes in human ovarian follicular fluid. *Fertil. Steril.* 1987; 47: 114–117.
13. **Lachapelle MH, Hemmings R, Roy DC, Falcone T, Miron P.** Flow cytometric evaluation of leukocyte subpopulations in the follicular fluids of infertile patients. *Fertil. Steril.* 1996; 65: 1135–1140.
14. **Lee A, Christenson LK, Patton PE, Burry KA, Stouffer RL.** Vascular endothelial growth factor production by human luteinized granulosa cells *in vitro*. *Hum. Reprod.* 1997; 12: 2756–2761.
15. **Loukides JA, Loy RA, Edwards R, Honig J, Visintin I, Polan ML.** Human follicular fluids contain tissue macrophages. *J. Clin. Endocrinol. Metab.* 1990; 71: 1363–1367.
16. **Otani N, Minami S, Yamoto M, Shikone T, Otani H, Nishiyama.** The vascular endothelial growth factor/fms-like tyrosine kinase system in human ovary during the menstrual cycle and early pregnancy. *J. Clin. Endocrinol. Metab.* 1999; 84: 3845–3851.
17. **Richards JS, Liu Z, Shimada M.** Immune-like mechanisms in ovulation. *Trends. End. Met.* 2008; 19: 191–196.
18. **Runesson E, Bostrom E, Janson PO, Brannstrom M.** The human preovulatory follicle is a source of the chemotactic cytokine interleukin-8. *Mol. Hum. Reprod.* 1996; 2: 245–250.
19. **Runesson E, Ivarsson K, Janson PO, Brannstrom M.** Gonadotropin- and cytokine-regulated expression of the chemokine interleukin-8 in the human preovulatory follicle of the menstrual cycle. *J. Clin. Endocrinol. Metab.* 2000; 85: 4387–4395.
20. **Scher AI, Petterson B, Blair E, Ellenberg JH, Grether JK, Haan E.** The risk of mortality or cerebral palsy in twins: a collaborative population-based study. *Pediatr. Res.* 2005; 52: 671–681.
21. **Sebire NJ, Jolly M, Harris J, Nicolaides KH, Regan L.** Risks of obstetric complications in multiple pregnancies: an analysis of more than 400,000 pregnancies in the UK. *Prenat. Neonatal. Med.* 2001, 6: 89–94.
22. **Smith MP, Flannery GR, Randle BJ, Jenkins JM, Holmes CH.** Leukocyte origin and profile in follicular aspirates at oocyte retrieval. *Hum Reprod.* 2005; 20:

- 3526–3531.
23. **Van Blerkom J, Antczak M, Schrader R.** The developmental potential of the human oocyte is related to the dissolved oxygen content of follicular fluid: association with vascular endothelial growth factor levels and perifollicular blood flow characteristics. *Hum. Reprod.* 1997; 12: 1047–1055.
  24. **Van Blerkom J.** Can the developmental competence of early human embryos be predicted effectively in the clinical IVF laboratory? *Hum. Reprod.* 1997; 12: 1610–1614.
  25. **Volarcik K, Sheean L, Goldfarb J, Woods L, Abdul-Karim FW, Hunt P.** The meiotic competence of *in-vitro* matured human oocytes is influenced by donor age: evidence that folliculogenesis is compromised in the reproductively aged ovary. *Hum. Reprod.* 1998; 13: 154–160.
  26. **World Health Organization.** WHO laboratory manual for the examination of human semen and sperm-cervical mucus interaction (4th ed), Cambridge Press, Cambridge, 1999, pp. 1–128.



Japan Society for Reproduction Engineering